

Causal Inference Methods and Case Studies

STAT24630

Jingshu Wang

Lecture 16

Topic: Assessing unconfoundedness, sensitivity analysis

- Assessing unconfoundedness
 - Negative control outcome
 - Negative control treatment
- Sensitivity analysis
 - Bound under no assumptions
 - Bound for the smoking example

Unconfoundedness and balance

- **Unconfoundedness property:** $W_i \perp (Y_i(0), Y_i(1)) \mid \mathbf{X}_i$
- This is an untestable assumption: we can never test for the unconfoundedness property as it is an assumption on the partially unmeasured potential outcomes

- We assess balancing of covariates and test for $W_i \perp \mathbf{X}_i \mid e(\mathbf{X}_i)$
- What we really care about is the balance of potential outcomes:

$$W_i \perp (Y_i(0), Y_i(1)) \mid e(\mathbf{X}_i)$$



within strata of observed covariates, potential outcomes corresponding to both treatment conditions need to be balanced between groups

- Covariate balancing is a necessary, but not sufficient condition, especially when there are unmeasured confounding pre-treatment covariates

Assessing unconfoundedness

- We can not test for unconfoundedness but we can assess the credibility of the unconfoundedness assumption indirectly
- Three approaches
 - **Negative control outcome:** choose proxy of the real outcome that
 1. Share a similar set of possible unmeasured confounding variables with the real outcome
 2. We know a priori that the treatment have **zero** causal effect on the proxy
 - **Negative control treatment:** choose new “treatment” that
 1. Share a similar set of possible unmeasured confounding variables with the real treatment
 2. We know a priori that the new “treatment” has **zero** causal effect on the outcome
 - **Assess robustness of the ATE estimation** given different sets of pre-treatment covariates

Negative control treatments and negative control outcomes



Negative control outcome (pseudo-outcome)

- One common way to find a good proxy of the outcome is the lagged outcome
 - E.x., outcome is the earning 1 year after treatment, lagged outcome is the earning 1 year before treatment
- The idea: the lagged outcome Y_i^{lag} , can be considered a proxy for $Y_i(0)$ and, given it is observed before the treatment, it is unaffected by the treatment
- By definition, the lagged outcome is also a pre-treatment covariate
 - Define $\mathbf{X}_i^r = \mathbf{X}_i \setminus Y_i^{lag}$, we test for the independence
$$H_0: W_i \perp Y_i^{lag} \mid \mathbf{X}_i^r$$
- In general, negative control outcome satisfies that $Y_i^{lag}(0) \equiv Y_i^{lag}(1)$, so we always observe its potential outcomes
- If we do not reject H_0 , it suggests that the unconfoundedness assumption is plausible.

Table 21.1. Summary Statistics for Selected Lottery Sample for the IRS Lottery Data

Variable	Label	All ($N = 496$)		Non-Winners ($N_t = 259$)	Winners ($N_c = 237$)	[t-Stat]	Nor Dif
		Mean	(S.D.)	Mean	Mean		
Year Won	(X_1)	6.23	(1.18)	6.38	6.06	-3.0	-0.27
Tickets Bought	(X_2)	3.33	(2.86)	2.19	4.57	9.9	0.90
Age	(X_3)	50.22	(13.68)	53.21	46.95	-5.2	-0.47
Male	(X_4)	0.63	(0.48)	0.67	0.58	-2.1	-0.19
Years of Schooling	(X_5)	13.73	(2.20)	14.43	12.97	-7.8	-0.70
Working Then	(X_6)	0.78	(0.41)	0.77	0.80	0.9	0.08
Earnings Year -6	(Y_{-6})	13.84	(13.36)	15.56	11.97	-3.0	-0.27
Earnings Year -5	(Y_{-5})	14.12	(13.76)	15.96	12.12	-3.2	-0.28
Earnings Year -4	(Y_{-4})	14.21	(14.06)	16.20	12.04	-3.4	-0.30
Earnings Year -3	(Y_{-3})	14.80	(14.77)	16.62	12.82	-2.9	-0.26
Earnings Year -2	(Y_{-2})	15.62	(15.27)	17.58	13.48	-3.0	-0.27
Earnings Year -1	(Y_{-1})	16.31	(15.70)	18.00	14.47	-2.5	-0.23
Pos Earnings Year -6	($Y_{-6} > 0$)	0.69	(0.46)	0.69	0.70	0.3	0.03
Pos Earnings Year -5	($Y_{-5} > 0$)	0.71	(0.45)	0.68	0.74	1.6	0.14
Pos Earnings Year -4	($Y_{-4} > 0$)	0.71	(0.45)	0.69	0.73	1.1	0.10
Pos Earnings Year -3	($Y_{-3} > 0$)	0.70	(0.46)	0.68	0.73	1.4	0.13
Pos Earnings Year -2	($Y_{-2} > 0$)	0.71	(0.46)	0.68	0.74	1.6	0.15
Pos Earnings Year -1	($Y_{-1} > 0$)	0.71	(0.45)	0.69	0.74	1.2	0.10

The
Imbens-
Rubin-
Sacerdote
lottery
data

The Imbens-Rubin-Sacerdote lottery data

Pseudo-Outcome	Remaining Covariates	Selected Covariates	Est	(s.e.)
Y_{-1}	$X_1, \dots, X_6, Y_{-6}, \dots, Y_{-2}, Y_{-6} > 0, \dots, Y_{-2} > 0$	X_2, X_5, X_6, Y_{-2}	-0.53	(0.58)
$\frac{Y_{-1}+Y_{-2}}{2}$	$X_1, \dots, X_6, Y_{-6}, \dots, Y_{-3}, Y_{-6} > 0, \dots, Y_{-3} > 0$	X_2, X_5, X_6, Y_{-3}	-1.16	(0.71)
$\frac{Y_{-1}+Y_{-2}+Y_{-3}}{3}$	$X_1, \dots, X_6, Y_{-6}, Y_{-5}, Y_{-4}, Y_{-6} > 0, Y_{-5} > 0, Y_{-4} > 0$	X_2, X_5, X_6, Y_{-4}	-0.39	(0.77)
$\frac{Y_{-1}+\dots+Y_{-4}}{4}$	$X_1, \dots, X_6, Y_{-6}, Y_{-5}, Y_{-6} > 0, Y_{-5} > 0$	X_2, X_5, X_6, Y_{-5}	-0.56	(0.89)
$\frac{Y_{-1}+\dots+Y_{-5}}{5}$	$X_1, \dots, X_6, Y_{-6}, Y_{-6} > 0$	X_2, X_5, X_6, Y_{-6}	-0.49	(0.87)
$\frac{Y_{-1}+\dots+Y_{-6}}{6}$	X_1, \dots, X_6	X_2, X_5, X_6	-2.56	(1.55)
Actual outcome Y	$X_1, \dots, X_6, Y_{-6}, \dots, Y_{-1}, Y_{-6} > 0, \dots, Y_{-1} > 0$	X_2, X_5, X_6, Y_{-1}	-5.74	(1.14)

Worse balance as no previous earnings are controlled

Negative control treatment (pseudo-treatment)

- One common case of negative control treatment is when there are multiple control groups
- Suppose we have two control groups and one treatment group $G_i \in \{c_1, c_2, t\}$ [e.g., ineligible, eligible nonparticipants and participants]

$$W_i = \begin{cases} 0 & \text{if } G_i = c_1, c_2, \\ 1 & \text{if } G_i = t. \end{cases}$$

- We test for

$$G_i \perp\!\!\!\perp Y_i(0) \mid X_i, G_i \in \{c_1, c_2\}$$

which is equivalent to

$$G_i \perp\!\!\!\perp Y_i^{\text{obs}} \mid X_i, G_i \in \{c_1, c_2\}.$$

Define pseudo-treatment for the lottery data

- One option is to have a comparison control group, of individuals who did not play the lottery at all
- Then we can compare between the “losers” and non-lottery players
- This comparison group is good because “losers” and non-lottery players can be substantially different due to various reasons (so they may share the same unmeasured confounders with that between “losers” and “winners”)
- However, we do not have such data
- Here, we split the winners into two subgroups
 - Median yearly prize for the winners is \$31,800
 - We treat the winners with yearly prize less than \$30,000 as the other group of control
 - Treat the winners with yearly prize larger than \$30,000 as the treated group

Pseudo-treatment analysis for the lottery data

Table 21.4. *Estimates of Average Difference in Outcomes for Controls and Small Winners (less than \$30,000) for the IRS Lottery Data*

Outcome	Subpopulation	Est	($\widehat{s. e.}$)
Y_i	All	-0.82	(1.37)
$\mathbf{1}_{Y_i=0}$	$Y_{i,-1} = 0$	-0.02	(0.05)
$\mathbf{1}_{Y_i=0}$	$Y_{i,-1} > 0$	0.07	(0.05)
Y_i	$Y_{i,-1} = 0$	-1.18	(1.10)
Y_i	$Y_{i,-1} > 0$	-0.16	(0.69)

Sensitivity analysis

- Most often, validity of unconfoundedness can not be easily checked. Alternatively, one should check sensitivity of a causal analysis to unconfoundedness
- **Sensitivity analysis** aims at assessing the bias of causal effect estimates when the unconfoundedness assumption is assumed to fail in some specific and meaningful ways
- Sensitivity is different from testing – unconfoundedness is intrinsically non-testable, more of a “insurance” check
- Sensitivity analysis in causal inference dates back to the Hill-Fisher debate on causation between smoking and lung cancer, and first formalized in Cornfield (1959, JNCI)

Bounds under no assumptions

- Consider a simple case where: 1. no covariates; 2. binary outcome
- We are interested in the ATE

$$\tau_{\text{sp}} = \mu_t - \mu_c,$$

where

$$\mu_t = \mathbb{E}[Y_i(1)] = p \cdot \mu_{t,1} + (1 - p) \cdot \mu_{t,0},$$

and

$$\mu_c = \mathbb{E}[Y_i(0)] = p \cdot \mu_{c,1} + (1 - p) \cdot \mu_{c,0}.$$

$$\mu_{t,1} = \mathbb{E}[Y_i(1)|W_i = 1]$$

$$\mu_{t,0} = \mathbb{E}[Y_i(1)|W_i = 0]$$

$$\mu_{c,1} = \mathbb{E}[Y_i(0)|W_i = 1]$$

$$\mu_{c,0} = \mathbb{E}[Y_i(0)|W_i = 0]$$

$$p = P(W_i = 1)$$

Identifiable
from
observed
data

Bound the unknown $\mu_{t,0}$ and $\mu_{c,1}$ by $[0, 1]$ as the outcome is binary

Bounds under no assumptions

- So we get the bounds

$$\begin{aligned}\mu_t &\in [p \cdot \mu_{t,1}, p \cdot \mu_{t,1} + (1 - p)] \\ \mu_c &\in [(1 - p) \cdot \mu_{c,0}, (1 - p) \cdot \mu_{c,0} + p]\end{aligned}$$

- The the bound of ATE $\tau = \tau_{sp} = \mu_t - \mu_c$ is

$$\tau \in [p \cdot \mu_{t,1} - (1 - p) \cdot \mu_{c,0} - p, p \cdot \mu_{t,1} + (1 - p) - (1 - p) \cdot \mu_{c,0}]$$

- Unfortunately, because we don't have any assumptions at all, this bound is not very informative:

we can easily show that $\tau^{upper} - \tau^{lower} \equiv 1$, so the bound always covers 0.

Result on the lottery data

- Binary outcome: whether the earning after treatment is positive or not
- Estimated quantities: $\hat{p} = \frac{N_t}{N} = 0.4675$, $\hat{\mu}_{t,1} = \bar{Y}_t^{\text{obs}} = 0.4106$ and $\hat{\mu}_{c,0} = \bar{Y}_c^{\text{obs}} = 0.5349$
- Plug in these quantities into our bound:
$$\tau \in [-0.56, 0.44]$$
- The two-sample difference estimate: $\bar{Y}_t^{\text{obs}} - \bar{Y}_c^{\text{obs}} = -0.134$


Sensitivity analysis bound: a more useful example

The smoking on lung cancer effect example (Cornfield et al. 1959 JNCI)

- Fisher argued the association between smoking and lung cancer may be due to a common gene that causes both
- Cornfield showed that if Fisher is right, we have $RR_{WU} \geq RR_{WY} \approx 9$
- Such a genetic confounder is too strong to be realistic
- Thus, smoking should have a causal effect on lung cancer

$$RR_{WY} = \frac{P[Y_i = 1|W_i = 1]}{P[Y_i = 1|W_i = 0]}$$

$$RR_{WU} = \frac{P[U_i = 1|W_i = 1]}{P[U_i = 1|W_i = 0]}$$



Sensitivity analysis bound: a more useful example

- Here, Y_i , U_i and W_i are all binary variables
- Define

$$p_0 = P[U_i = 1|W_i = 0], \quad p_1 = P[U_i = 1|W_i = 1]$$

- If there is no causal effect of smoking on lung cancer, then

$$P[Y_i = 1|W_i = 0, U_i = 0] = P[Y_i = 1|W_i = 1, U_i = 0] = r_0,$$

$$P[Y_i = 1|W_i = 0, U_i = 1] = P[Y_i = 1|W_i = 1, U_i = 1] = r_1$$

- Then we have

$$RR_{AY} = \frac{P[Y_i = 1|W_i = 1]}{P[Y_i = 1|W_i = 0]} = \frac{r_0(1 - p_1) + r_1p_1}{r_0(1 - p_0) + r_1p_0}$$

- Let $p_1 \geq p_0$, then because we observe $RR_{AY} > 1$, then (from some math)

$$RR_{WY} = \frac{r_0(1 - p_1) + r_1p_1}{r_0(1 - p_0) + r_1p_0} \leq \frac{p_1}{p_0} = RR_{WU}$$