

STAT347: Generalized Linear Models

Lecture 2

Winter, 2024
Jingshu Wang

Today's topics:

- The exponential dispersion family
- Exponential family distribution for GLM
- Likelihood score equations for parameter estimation
- Reading: Agresti Chapters 4.1-4.2, Faraway Chapter 8.1-8.2

The exponential dispersion family

- A random variable Y follows an exponential dispersion family distribution and has the density $f(y; \theta, \phi)$ of the form

$$f(y; \theta, \phi) = e^{\frac{y\theta - b(\theta)}{a(\phi)}} f_0(y; \phi)$$

Terminologies:

- θ : natural or canonical parameters
 - $b(\theta)$: normalizing or cumulant function
 - ϕ : dispersion parameter with $a(\phi) > 0$
 - Typically $a(\phi) \equiv 1$ and $f_0(y; \phi) = f_0(y)$. An exception is the Gaussian distribution where $a(\phi) = \sigma^2$
-
- “density” here includes the possibility of discrete atoms.
 - Above definition is not the most general form of the exponential family distribution

Some well-known examples

- Normal distribution for continuous data

$$f(y; \mu, \sigma) = e^{\frac{y\mu - \mu^2/2}{\sigma^2}} \left[\frac{1}{\sqrt{2\pi\sigma}} e^{-\frac{y^2}{2\sigma^2}} \right]$$

- $\theta = \mu, b(\theta) = \theta^2/2, a(\phi) = \sigma^2$

- Bernoulli distribution for binary data

$$\begin{aligned} f(y; p) &= p^y (1-p)^{1-y} = e^{y \log \frac{p}{1-p} + \log(1-p)} \\ &= e^{y\theta - \log[1+e^\theta]} \end{aligned}$$

- $\theta = \log\left(\frac{p}{1-p}\right), b(\theta) = \log[1 + e^\theta], a(\phi) = 1$

Some well-known examples

- Binomial distribution for counts data

$$\begin{aligned} f(y; p, n) &= \binom{n}{y} p^y (1-p)^{n-y} = e^{y \log \frac{p}{1-p} + n \log(1-p)} \binom{n}{y} \\ &= e^{y\theta - n \log[1+e^\theta]} \binom{n}{y} \end{aligned}$$

- $\theta = \log\left(\frac{p}{1-p}\right)$, $b(\theta) = n \log[1 + e^\theta]$, $a(\phi) = 1$

- Poisson distribution for counts data

$$f(y; \lambda) = \frac{e^{-\lambda} \lambda^y}{y!} = e^{y \log \lambda - \lambda} \frac{1}{y!} = e^{y\theta - e^\theta} \frac{1}{y!}$$


- $\theta = \log(\lambda)$, $b(\theta) = e^\theta$, $a(\phi) = 1$

Some well-known examples

- Gamma distribution for positive real-valued data

$$f(y; k, \theta) = \frac{1}{\Gamma(k)\theta^k} y^{k-1} e^{-y/\theta}$$
$$= e^{\frac{-\frac{1}{k\theta}y + \log\left(\frac{1}{k\theta}\right)}{1/k}} \frac{y^{k-1} k^k}{\Gamma(k)}$$

- Mean $\mu = k\theta$, variance $k\theta^2 = \mu^2/k$
- Canonical parameter $-1/\mu$, dispersion parameter $1/k$



Exponential family distribution for GLM

- Assume that each observation y_i follows an exponential family with the canonical parameter θ_i and a shared dispersion parameter ϕ
- $\mu_i = \mathbb{E}(y_i)$ is a function of X_i , so θ_i is also a function of X_i
- Canonical link function

$$g(\mu_i) = \theta_i = X_i^T \beta$$

Canonical link function examples

- Gaussian: $\theta_i = \mu_i = X_i^T \beta$
- Binomial and Bernoulli distribution: $\theta_i = \log\left(\frac{p_i}{1-p_i}\right) = X_i^T \beta$
 - Called the logit function
- Poisson distribution: $\theta_i = \log(\mu_i) = X_i^T \beta$

Moment relationships

- The exponential family has some special properties that can make our calculation easier
 - Calculate mean and variance of Y

$$\mu = \mathbb{E}(y) = b'(\theta)$$

$$V_{\theta} = \text{Var}(y) = b''(\theta)a(\phi)$$

- Why? As $\int f(y; \theta, \phi) dy = 1$, we have

$$e^{b(\theta)/a(\phi)} = \int e^{y\theta/a(\phi)} f_0(y; \phi) dy$$

- Take derivatives with respect to θ

Moment relationships

- The exponential family has some special properties that can make our calculation easier
 - Calculate mean and variance of Y

$$\mu = \mathbb{E}(Y) = b'(\theta)$$

$$V_{\theta} = \text{Var}(Y) = b''(\theta)a(\phi)$$

- The above relationship also indicates that

$$\frac{\partial \mu}{\partial \theta} = \frac{\text{Var}(Y)}{a(\phi)} > 0$$

- Mapping from θ to μ is one to one increasing

Likelihood score equations

- We now use the maximum likelihood method to solve for the GLM and estimate β

Assume each observation y_i follows an exponential dispersion distribution

$$f(y_i; \theta_i, \phi) = e^{\frac{y_i \theta_i - b(\theta_i)}{a(\phi)}} f_0(y_i; \phi)$$

and the link function $g(\mu_i) = X_i^T \beta$. Then for n independent observations, the log likelihood is

$$L = \sum_i L_i = \sum_i \frac{y_i \theta_i - b(\theta_i)}{a(\phi)} + \sum_i \log f_0(y_i; \phi)$$

Likelihood score equation for the canonical link

If $g(\mu_i) = \theta_i = X_i^T \beta$, then

$$L = \frac{1}{a(\phi)} \left[\sum_j \left(\sum_i y_i x_{ij} \right) \beta_j - \sum_i b(X_i^T \beta) \right] + \sum_i \log f_0(y_i; \phi)$$

- Score equation for β_j

$$\frac{\partial L}{\partial \beta_j} = \frac{1}{a(\phi)} \left[\sum_i y_i x_{ij} - \sum_i b'(X_i^T \beta) x_{ij} \right] = \frac{1}{a(\phi)} \left[\sum_i (y_i - \mu_i) x_{ij} \right] = 0$$

which is equivalent to

$$\sum_i (y_i - \mu_i) x_{ij} = 0$$

Likelihood score equation for the canonical link

- Examples

Gaussian model:

$$\sum_i (y_i - X_i^T \beta) x_{ij} = 0$$

Poisson model:

$$\sum_i (y_i - e^{X_i^T \beta}) x_{ij} = 0$$

- L is a concave function of $\beta = (\beta_1, \dots, \beta_p)$

$$\frac{\partial}{\partial \beta} \left[\sum_i (y_i - \mu_i) X_i \right] = - \sum_i \frac{\partial \mu_i}{\partial \theta_i} \frac{\partial \theta_i}{\partial \beta} X_i^T = - \sum_i \frac{\text{Var}(y_i)}{a(\phi)} X_i X_i^T \prec 0$$

- Easy optimization to find the solution (will discuss computation later)

Likelihood score equation for a general link

Let $\eta_i = g(\mu_i) = X_i^T \beta$ Then

$$\frac{\partial L_i}{\partial \beta_j} = \frac{\partial L_i}{\partial \theta_i} \frac{\partial \theta_i}{\partial \mu_i} \frac{\partial \mu_i}{\partial \eta_i} \frac{\partial \eta_i}{\partial \beta_j}$$

We have

- $\frac{\partial L_i}{\partial \theta_i} = \frac{y_i - b'(\theta_i)}{a(\phi)} = \frac{y_i - \mu_i}{a(\phi)}$
- $\frac{\partial \theta_i}{\partial \mu_i} = \frac{1}{b''(\theta_i)} = \frac{a(\phi)}{\text{Var}(y_i)}$
- $\frac{\partial \mu_i}{\partial \eta_i} = \frac{\partial \mu_i}{\partial g(\mu_i)} = \frac{1}{g'(\mu_i)}$
- $\frac{\partial \eta_i}{\partial \beta_j} = x_{ij}$

Likelihood score equation for a general link

- The score equations can be written as

$$\frac{\partial L}{\partial \beta_j} = \sum_i \frac{(y_i - \mu_i) x_{ij}}{\text{Var}(y_i)} \frac{1}{g'(\mu_i)} = 0$$

- μ_i and $\text{Var}(y_i)$ are both functions of $\beta = (\beta_1, \dots, \beta_p)$
- The score equations only depend on the mean and variance of y_i
- Matrix form of the score equation:

$$\dot{L}(\beta) = X^T D V^{-1} (y - \mu) = 0$$

where $V = \text{diag}(\text{Var}(y_1), \dots, \text{Var}(y_n))$ and $D = \text{diag}(g'(\mu_1), \dots, g'(\mu_n))^{-1}$,
 $y = (y_1, \dots, y_n)$ and $\mu = (\mu_1, \dots, \mu_n)$.

- L is not necessarily a concave function of β

Likelihood score equation for a general link

Special cases

- If the link function is the canonical link, then $D = \frac{1}{a(\phi)}V$, thus the score equation becomes

$$\frac{1}{a(\phi)}X^T(y - \mu) = 0$$

the same as we derived earlier

- If we assume that $g(\mu_i) = \mu_i = X_i^T \beta$, then the estimating (score) equation becomes

$$\sum_i \frac{(y_i - X_i^T \beta) X_i}{\text{Var}(y_i)} = 0$$

which looks like weighted least square (difference: weights can depend on β)